

Antibiotic Resistance Awareness


AMR Worldwide

14,000 Patients Die

of *C.difficile* infection annually in the **USA**.⁽¹⁾ The use of antibiotics was a major contributing factor in up to 85% of cases.⁽²⁾



23,000 Patients Die Each Year as a result of **antibiotic-resistant infections** in the **USA**.⁽¹⁾



2,000,000 Infections per year contain bacteria that are resistant to one or more antibiotics in the **USA**.⁽¹⁾



11,000 Estimated Deaths caused by methicillin-resistant Staphylococcus aureus (**MRSA**) each year in the **USA**.⁽⁵⁾



25,000 Patients Die Each Year as a result of antibiotic-resistant infections in **Europe**.⁽³⁾



400,000 Infections per year with the 6 most frequent multi-drug resistant (MDR) bacteria, in 4 types of infection, in **Europe**.⁽⁶⁾



480,000 People Infected by drug-resistant TB strains in 2013 **Worldwide**.⁽⁴⁾



1 Child Dies Every 9 Minutes from an infection caused by antibiotic-resistant bacteria in **India**.⁽⁷⁾



(1) Antibiotic Resistance Threats in the United States, 2013. Centers for Disease Control and Prevention (CDC)

(2) Chang HT et al. Infect Control Hosp Epidemiol, 2007; 28:926-931

(3) ENVI 2013-04-10 European Report on patient safety

(4) Global Tuberculosis Report. World Health Organization, 2014

(5) WHO Antimicrobial Resistance Fact sheet N°194. Updated April 2015.

(6) Antibiotic Resistance: Implications for Global Health and Novel Intervention Strategies. Washington, DC. The National Academies Press. 2010 (Source ECDC/ EMEA 2009)

(7) Laxminarayan et al. Antibiotic resistance - the need for global solutions. Lancet Infect Dis 2013;13:1057-98